Fluid Phase Equilibria, 44 (1989) 365–366
Elsevier Science Publishers B.V., Amsterdam – Printed in The Netherlands

Contents Volume 44 (1988/1989)

Volume 44, No. 1

A flexible representation of quaternary fluid phase equilibria for use in a chemical flood simulator
R.W.S. Foulser and J.A. Oakes (Dorchester, Gt. Britain) 1–21
Analysis of retrograde behavior and the cross-over effect in supercritical fluids
E.H. Chimowitz, F.D. Kelley and F.M. Munoz (Rochester, NY, U.S.A.) 23–52
Measurement and prediction of liquid–liquid equilibria of ethyl acetate–methanol–water in the presence of dissolved inorganic salts
M. Rajendran and D. Srinivasan (Madras, India) 53–75
Vapour–liquid equilibria. IV. The ternary system carbon tetrachloride–methanol–chloroform at 293.15 K
M. Góral, P. Oracz and S. Warycha (Warsaw, Poland) 77–93
Phase equilibria in binary mixtures of propane + acenaphthene
C.J. Peters, J.L. de Roo and J. de Swaan Arons (Delft, The Netherlands) 95–103
Solubilities of carbon dioxide and hydrogen sulfide in propylene carbonate, *N*-methylpyrrolidone and sulfolane
F. Murrieta-Guevara, A. Romero-Martinez and A. Trejo (Mexico, Mexico) . . 105–115
Ternary phase equilibrium studies of the water–ethanol–1,8-cineole system
A.F.M. Barton and J. Tjandra (Perth, Australia) 117–123

Volume 44, No. 2

Prediction of mixed gas solubility at high pressure
A.K. Myers (Princeton, NJ, U.S.A.) and A.L. Myers (Philadelphia, PA, U.S.A.) 125–144
Group contribution methods for molecular mixtures. II. Computer simulation results
C. Massobrio and J.M. Haile (Clemson, SC, U.S.A.) and L.L. Lee (Norman, OK, U.S.A.) 145–173
Group contribution methods for molecular mixtures. III. Solution of groups tested via computer simulation
J.M. Haile (Clemson, SC, U.S.A.) and L.L. Lee (Norman, OK, U.S.A.) 175–194
Binary vapor–liquid–liquid equilibrium of dimethyl ether–water and mutual solubilities of methyl chloride and water: experimental results and data reduction
H. Holldorff and H. Knapp (Berlin, F.R.G.) 195–209
Vapour pressure of binary, three-phase (S–L–V) systems and solubility
A. Ksiązczak and J.J. Kosinski (Warsaw, Poland) 211–236

Volume 44, No. 3

Phase equilibria in model binary mixtures. I. Dipolar hard spheres of equal size
W.R. Smith (Guelph, Ontario, Canada), O.H. Scalise (La Plata, Argentina) and T.W. Melnyk (Pinawa, Manitoba, Canada) 237–254
Prediction of vapor–liquid equilibria in binary mixtures containing polar components from an extended Lee–Kesler corresponding-states technique
J.K. Johnson and R.L. Rowley (Provo, UT, U.S.A.) 255–272
Note on the performance of the quasi-chemical group solution theory
J.A. Abusleme and J.H. Vera (Montreal, Quebec, Canada) 273–283
Excess volumes of cyclohexane + a linear ether at 298.15 K
P. Berti, L. Lepori and E. Matteoli (Pisa, Italy) 285–294
Observations on the multiphase equilibria behavior of CO_2-rich and ethane-rich mixtures
M.M. Miller and K.D. Luks (Tulsa, OK, U.S.A.) 295–304
A comparison of solid–liquid equilibrium with vapor–liquid equilibrium for prediction of activity coefficients in systems containing polynuclear aromatics
J.E. Coon, J.E. Auwaerter and E. McLaughlin (Baton Rouge, LA, U.S.A.) . . . 305–345
Thermodynamics of methylenebromide + aromatic hydrocarbon mixtures: molar excess Gibbs free energy of mixing
P.P. Singh and V.K. Sharma (Rohtak, India) 347–355

Short Communication

Excess volumes of binary mixtures of trichloroethylene with some aliphatic, alicyclic and aromatic hydrocarbons
K. Ramanjaneyulu, K.N. Surendranath and A. Krishnaiah (Tirupati, India) . 357–361

Author Index 363

Contents of Volume 44 (1988/1989) 365–366